


Manufacturer/Supplier:				
WUXI DONGLIN SCI&TECH DEVELOPMENT CO., LTD				
Address: A1-203 Mingpin CityII,No.8 Xihudong Road.Liangxi District				
Wuxi Jiangsu Province, China				
TELE: 86-510-82732223		FAX:	86-510-82720101-8014	
Web : <u>www.dldevelop.com</u>				
Email: <u>info@dldevelop.com.cn</u> <u>service@dldevelop.com.cn</u>				
Product name:	Rabbit anti-Transgelin-2 Antibody			
Catalog:	DL96333A			
Synonyms:	KIAA0120; Transgelin-2; Epididymis tissue protein			
	Li 7e; SM22-alp	oha homolo	g	
_				
Immunogen:	Recombinant full length protein of human			
	Transgelin-2			
Form: Liquid		Concentr	ation: 1mg/mL	
Form. Elquid		Concenti		
Size: 100 ul/50 ul		Host: Rat	bit	
		110.500 1100		
Reactivity: Human, Mouse, Rat		Applicati	on: WB, IF/IC	
Clonality: Polyclonal		Dilution: WB (1/500 - 1/2000),		
		IF/IC (1/5	0 - 1/200)	



Entrez Gene:	8407/21346/304983
SwissProt:	P37802/Q9WVA4/Q5XFX0
Purification:	The antibody was purified by immunogen affinity chromatography.
Buffer:	Liquid in 0.42% Potassium phosphate, 0.87% Sodium chloride, pH 7.3, 30% glycerol, and 0.01% sodium azide.

WB description:


Western blot analysis of Transgelin-2 expression in HEK293T (A), Hela (B), THP1 (C), MCF7 (D), SKOV3 (E), mouse lung (F) whole cell lysates.



IF/ICC description:



Immunofluorescent analysis of Transgelin-2 staining in MCF7 cells. Formalin-fixed cells were permeabilized with 0.1% Triton X-100 in TBS for 5-10 minutes and blocked with 3% BSA-PBS for 30 minutes at room temperature. Cells were probed with the primary an



Storage:

Store at -20°C. Avoid repeated freeze / thaw cycles.

