

Manufacturer/Supplier:

WUXI DONGLIN SCI&TECH DEVELOPMENT CO., LTD

Address: A1-203 Mingpin CityII, No.8 Xihudong Road, Liangxi District
Wuxi Jiangsu Province, China

TELE: 86-510-82732223

FAX: 86-510-82720101-8014

Web : www.dldevelop.com

Email: info@dldevelop.com.cn service@dldevelop.com.cn

Product name: Rabbit anti-DLST Antibody

Catalog: DL96351A

Synonyms: DLTS; Dihydrolipoyllysine-residue
succinyltransferase component of 2-oxoglutarate
dehydrogenase complex, mitochondrial;
2-oxoglutarate dehydrogenase complex component
E2; OGDC-E2; Dihydrolipoamide
succinyltransferase component of 2-oxoglutarate
dehydrogen

Immunogen: Recombinant full length protein of human DLST

Form: Liquid

Concentration: 1mg/mL

Size: 100 ul/50 ul

Host: Rabbit

Reactivity: Human, Mouse, Rat

Application: WB, IHC

Clonality: Polyclonal

Dilution: WB (1/500 - 1/2000),
IHC (1/50 - 1/200)

Entrez Gene: 1743/78920/299201


SwissProt: P36957/Q9D2G2/Q01205

Purification: The antibody was purified by immunogen affinity chromatography.

Buffer: Liquid in 0.42% Potassium phosphate, 0.87% Sodium chloride, pH 7.3, 30% glycerol, and 0.01% sodium azide.

WB description:

Western blot analysis of DLST expression in SKOV3 (A), Hela (B), mouse kidney (C), rat liver (D) whole cell lysates.



IHC description:

Immunohistochemical analysis of DLST staining in human liver cancer formalin fixed paraffin embedded tissue section. The section was pre-treated using heat mediated antigen retrieval with sodium citrate buffer (pH 6.0). The section was then incubated with the antibody at room temperature and detected using an HRP conjugated compact polymer system. DAB was used as the chromogen. The section was then counterstained with haematoxylin and mounted with DPX.

Storage:

Store at -20°C. Avoid repeated freeze / thaw cycles.