


Manufacturer/Supplier:				
WUXI DONGLIN SCI&TECH DEVELOPMENT CO., LTD				
Address: A1-203 Mingpin CityII,No.8 Xihudong Road.Liangxi District				
Wuxi Jiangsu Province, China				
TELE: 86-510-82	2732223	FAX:	86-510-82720101-8014	
Web : <u>www.dldevelop.com</u>				
Email: <u>info@dldevelop.com.cn</u> <u>service@dldevelop.com.cn</u>				
Product name: Catalog:	Rabbit anti-eIF3 DL96379A	C Antibod	ÿ	
Synonyms:	EIF3S8; Eukaryotic translation initiation factor 3 subunit C; eIF3c; Eukaryotic translation initiation factor 3 subunit 8; eIF3 p110			
Immunogen:	Recombinant full length protein of human eIF3C			
Form: Liquid		Concentr	ation: 1mg/mL	
Size: 100 ul/50 ul		Host: Rabbit		
Reactivity: Human, Mouse, Rat		Application: WB, IF/IC		
Clonality: Polyclonal		Dilution: WB (1/500 - 1/2000), IF/IC (1/50 - 1/100)		

DEVELOP.

Entrez Gene:	8663/56347/293484
SwissProt:	Q99613/Q8R1B4/B5DFC8
Purification:	The antibody was purified by immunogen affinity chromatography.
Buffer:	Liquid in 0.42% Potassium phosphate, 0.87% Sodium chloride, pH 7.3, 30% glycerol, and 0.01% sodium azide.

WB description:

Western blot analysis of eIF3C expression in HepG2 (A), SW620 (B), mouse testis (C) whole cell lysates.




IF/ICC description:

Immunofluorescent analysis of eIF3C staining in U2OS cells.



Formalin-fixed cells were permeabilized with 0.1% Triton X-100 in TBS for 5-10 minutes and blocked with 3% BSA-PBS for 30 minutes at room temperature. Cells were probed with the primary antibody



Storage:

Store at -20°C. Avoid repeated freeze / thaw cycles.

