

Manufacturer/Supplier:

WUXI DONGLIN SCI&TECH DEVELOPMENT CO., LTD

Address: A1-203 Mingpin CityII, No.8 Xihudong Road. Liangxi District

Wuxi Jiangsu Province, China

TELE: 86-510-82732223 FAX: 86-510-82720101-8014

Web: www.dldevelop.com

Email: info@dldevelop.com.cn service@dldevelop.com.cn

Product name: Rabbit anti-eIF3H Antibody

Catalog: DL96380A

EIF3S3; Eukaryotic translation initiation factor 3 **Synonyms:**

> subunit H; eIF3h; Eukaryotic translation initiation factor 3 subunit 3; eIF-3-gamma; eIF3 p40 subunit

Immunogen: Recombinant full length protein of human eIF3H

Form: Liquid Concentration: 1mg/mL

Size: 100 ul/50 ul Host: Rabbit

Reactivity: Human, Mouse, Rat **Application:** WB, IF/IC

Clonality: Polyclonal **Dilution:** WB (1/500 - 1/2000),

IF/IC (1/50 - 1/100)



Entrez Gene: 8667/68135/100911110; 299899

O15372/O91WK2/O6P9U8 SwissProt:

Purification: The antibody was purified by immunogen affinity

chromatography.


Buffer: Liquid in 0.42% Potassium phosphate, 0.87%

Sodium chloride, pH 7.3, 30% glycerol, and 0.01%

sodium azide.

WB description:

Western blot analysis of eIF3H expression in HL60 (A), HT29 (B), NIH3T3 (C), mouse testis (D), rat spleen (E) whole cell lysates.



IF/ICC description:

Immunofluorescent analysis of eIF3H staining in U2OS cells.



Formalin-fixed cells were permeabilized with 0.1% Triton X-100 in TBS for 5-10 minutes and blocked with 3% BSA-PBS for 30 minutes at room temperature. Cells were probed with the primary antibody



Storage:

Store at -20°C. Avoid repeated freeze / thaw cycles.