


M f 4 /S				
Manufacturer/Supplier:				
WUXI DONGLIN SCI&TECH DEVELOPMENT CO., LTD				
Address: A1-203 Mingpin CityII,No.8 Xihudong Road.Liangxi District				
Wuxi Jiangsu Province, China				
TELE: 86-510-82732223		FAX:	86-510-82720101-8014	
Web : <u>www.dldevelop.com</u>				
Email: info@dldevelop.com.cn service@dldevelop.com.cn				
Product name:	Rabbit anti-Aspartoacylase Antibody			
Catalog:	DL96421A			
Catalog.	DLJ0421A			
C				
Synonyms:	ACY2; ASP; Aspartoacylase; Aminoacylase-2;			
	ACY-2			
Immunogen:	Recombinant fu	ll length pro	otein of human	
	Aspartoacylase			
Form: Liquid		Concentr	ation: 1mg/mL	
Size: 100 ul/50 ul		Host: Rab	bit	
		110000 11000		
Reactivity: Human		Applicati	on: WB, IF/IC	
Reactivity. Human		Аррисан	on. wb, n/iC	
Clanality Dalualanal		Dilution	WD(1/500, 1/2000)	
Clonality: Polyclonal		Dilution: WB (1/500 - 1/2000),		
		IF/IC (1/5	0 - 1/100)	

Entrez Gene:	443
SwissProt:	P45381
Purification:	The antibody was purified by immunogen affinity chromatography.
Buffer:	Liquid in 0.42% Potassium phosphate, 0.87% Sodium chloride, pH 7.3, 30% glycerol, and 0.01% sodium azide.

WB description:

Western blot analysis of Aspartoacylase expression in Hela (A), mouse kidney (B), rat kidney (C) whole cell lysates.




IF/ICC description:

Immunofluorescent analysis of Aspartoacylase staining in MCF7



cells. Formalin-fixed cells were permeabilized with 0.1% Triton X-100 in TBS for 5-10 minutes and blocked with 3% BSA-PBS for 30 minutes at room temperature. Cells were probed with the primary



Storage:

Store at -20°C. Avoid repeated freeze / thaw cycles.

