

Manufacturer/Supplier:				
WUXI DONGLIN SCI&TECH DEVELOPMENT CO., LTD				
Address: A1-203 Mingpin CityII,No.8 Xihudong Road.Liangxi District				
Wuxi Jiangsu Province, China				
TELE: 86-510-82	2732223	FAX:	86-510-82720101-8014	
Web : <u>www.dldevelop.com</u>				
Email: info@dldevelop.com.cn service@dldevelop.com.cn				
Product name:	Rabbit anti-AUH Antibody			
Catalog:	DL96456A			
8				
Synonyms:	Methylglutaconyl-CoA hydratase, mitochondrial;			
~ j == 0 == j === 0 0	AU-specific RNA-binding enoyl-CoA hydratase;			
	AU-binding pro	U		
	re-omaing pro	tern/enoyi-v	cont injuratase	
Immunogen:	Recombinant full length protein of human AUH			
ininunogen.	Recombinant Iu	n lengti pr		
Form: Liquid		Concentr	ation: lmg/mL	
Form. Elquid		Concentr	ation. Ting/InL	
Size: 100 ul/50 ul		Host: Rat	bit	
Size. 100 ul/30 ul		Host. Ruc		
Reactivity: Human, Mouse, Rat		Annlicati	on: WB, IHC	
reactivity. Human, mouse, Rat		rippiicati		
Clonality: Polyclonal		Dilution:	WB (1/500 - 1/2000),	
		IHC (1/50 - 1/200)		
		110 (1/50	1/200)	


DEVELOP.

Entrez Gene:	549/11992
SwissProt:	Q13825/Q9JLZ3
Purification:	The antibody was purified by immunogen affinity chromatography.
Buffer:	Liquid in 0.42% Potassium phosphate, 0.87% Sodium chloride, pH 7.3, 30% glycerol, and 0.01% sodium azide.

WB description:

Western blot analysis of AUH expression in A549 (A), SW480

(B), mouse brain (C), mouse kidney (D), rat heart (E) whole cell lysates.



IHC description:

Immunohistochemical analysis of AUH staining in rat kidney



formalin fixed paraffin embedded tissue section. The section was pre-treated using heat mediated antigen retrieval with sodium citrate buffer (pH 6.0). The section was then incubated with the antibody at room temperature and detected using an HRP conjugated compact polymer system. DAB was used as the chromogen. The section was then counterstained with haematoxylin and mounted with DPX.

Storage:

Store at -20°C. Avoid repeated freeze / thaw cycles.