

Manufacturer/Supplier:

WUXI DONGLIN SCI&TECH DEVELOPMENT CO., LTD

Address: A1-203 Mingpin CityII, No.8 Xihudong Road. Liangxi District

Wuxi Jiangsu Province, China

TELE: 86-510-82732223 FAX: 86-510-82720101-8014

Web: www.dldevelop.com

Email: info@dldevelop.com.cn service@dldevelop.com.cn

Product name: Rabbit anti-FURIN Antibody

Catalog: DL96831A

FUR; PACE; PCSK3; Furin; Dibasic-processing **Synonyms:**

enzyme; Paired basic amino acid residue-cleaving

enzyme; PACE

Immunogen: Recombinant full length protein of human FURIN

Form: Liquid Concentration: 1mg/mL

Size: 100 ul/50 ul Host: Rabbit

Reactivity: Human, Mouse, Rat **Application:** WB, IHC

Clonality: Polyclonal **Dilution:** WB (1/500 - 1/1000),

IHC (1/50 - 1/200)



Entrez Gene: 5045/18550/54281

P09958/P23188/P23377 SwissProt:

Purification: The antibody was purified by immunogen affinity

chromatography.


Buffer: Liquid in 0.42% Potassium phosphate, 0.87%

Sodium chloride, pH 7.3, 30% glycerol, and 0.01%

sodium azide.

WB description:

Western blot analysis of FURIN expression in A375 (A), mouse ovary (B) whole cell lysates.



IHC description:

Immunohistochemical analysis of FURIN staining in human liver



cancer formalin fixed paraffin embedded tissue section. The section was pre-treated using heat mediated antigen retrieval with sodium citrate buffer (pH 6.0). The section was then incubated with the antibody at room temperature and detected using an HRP conjugated compact polymer system. DAB was used as the chromogen. The section was then counterstained with haematoxylin and mounted with DPX.

Storage:

Store at -20°C. Avoid repeated freeze / thaw cycles.