

**Manufacturer/Supplier:**

WUXI DONGLIN SCI&TECH DEVELOPMENT CO., LTD

**Address:** A1-203 Mingpin CityII, No.8 Xihudong Road, Liangxi District  
Wuxi Jiangsu Province, China

TELE: 86-510-82732223

FAX: 86-510-82720101-8014

**Web :** [www.dldevelop.com](http://www.dldevelop.com)

Email: [info@dldevelop.com.cn](mailto:info@dldevelop.com.cn) [service@dldevelop.com.cn](mailto:service@dldevelop.com.cn)

**Product name:** Rabbit anti-GLUR3 Antibody

**Catalog:** DL96851A

**Synonyms:** GLUR3; GLURC; Glutamate receptor 3; GluR-3;  
AMPA-selective glutamate receptor 3; GluR-C;  
GluR-K3; Glutamate receptor ionotropic, AMPA 3;  
GluA3

**Immunogen:** Recombinant full length protein of human GLUR3

**Form:** Liquid

**Concentration:** 1mg/mL

**Size:** 100 ul/50 ul

**Host:** Rabbit

**Reactivity:** Human, Mouse, Rat

**Application:** WB, IF/IC

**Clonality:** Polyclonal

**Dilution:** WB (1/500 - 1/2000),

IF/IC (1/50 - 1/200)

**Entrez Gene:** 2892/53623/29628


**SwissProt:** P42263/Q9Z2W9/P19492

**Purification:** The antibody was purified by immunogen affinity chromatography.

**Buffer:** Liquid in 0.42% Potassium phosphate, 0.87% Sodium chloride, pH 7.3, 30% glycerol, and 0.01% sodium azide.


### WB description:

Western blot analysis of GLUR3 expression in U251 (A), PC3 (B), mouse brain (C), mouse lung (D) whole cell lysates.



### IF/ICC description:

Immunofluorescent analysis of GLUR3 staining in U2OS cells. Formalin-fixed cells were permeabilized with 0.1% Triton X-100 in TBS for 5-10 minutes and blocked with 3% BSA-PBS for 30 minutes at room temperature. Cells were probed with the primary antibody



**Storage:**

Store at -20°C. Avoid repeated freeze / thaw cycles.