

Manufacturer/Supplier:				
WUXI DONGLIN SCI&TECH DEVELOPMENT CO., LTD				
Address: A1-203 Mingpin CityII,No.8 Xihudong Road.Liangxi District				
Wuxi Jiangsu Province, China				
TELE: 86-510-82732223		FAX:	86-510-82720101-8014	
Web : <u>www.dldevelop.com</u>				
Email: <u>info@dldevelop.com.cn</u> <u>service@dldevelop.com.cn</u>				
Product name: Catalog:	Rabbit anti-IGF2 DL96886A	BP3 Antib	ody	
Synonyms:	IMP3; KOC1; VICKZ3; Insulin-like growth factor 2 mRNA-binding protein 3; IGF2 mRNA-binding protein 3; IMP-3; IGF-II mRNA-binding protein 3; KH domain-containing protein overexpressed in cancer; hKOC; VICKZ family member 3			
Immunogen:	Recombinant ful	l length pro	otein of human IGF2BP3	
Form: Liquid		Concentra	ation: 1mg/mL	
Size: 100 ul/50 ul		Host: Rab	bit	
Reactivity: Human		Application: WB, IHC		



Clonality: Polyclonal


Dilution: WB (1/500 - 1/1000), IHC (1/50 - 1/200)

Entrez Gene:	10643
SwissProt:	O00425
Purification:	The antibody was purified by immunogen affinity chromatography.
Buffer:	Liquid in 0.42% Potassium phosphate, 0.87% Sodium chloride, pH 7.3, 30% glycerol, and 0.01% sodium azide.

WB description:

Western blot analysis of IGF2BP3 expression in Hela (A), PC3 (B), NIH3T3 (C) whole cell lysates.

DEVELOP



IHC description:

Immunohistochemical analysis of IGF2BP3 staining in mouse brain formalin fixed paraffin embedded tissue section. The section was pre-treated using heat mediated antigen retrieval with sodium citrate buffer (pH 6.0). The section was then incubated with the antibody at room temperature and detected using an HRP conjugated compact polymer system. DAB was used as the chromogen. The section was then counterstained with haematoxylin and mounted with DPX.

Storage:

Store at -20°C. Avoid repeated freeze / thaw cycles.