

Manufacturer/Supplier:				
WUXI DONGLIN SCI&TECH DEVELOPMENT CO., LTD				
Address: A1-203 Mingpin CityII,No.8 Xihudong Road.Liangxi District				
Wuxi Jiangsu Province, China				
TELE: 86-510-82732223 FAX: 86-51			86-510-82720101-8014	
Web : <u>www.dldevelop.com</u>				
Email: <u>info@dldevelop.com.cn</u> <u>service@dldevelop.com.cn</u>				
Product name:	Rabbit anti-NDUFS4 Antibody			
Catalog:	DL96967A			
Synonyms:	NADH dehydrogenase [ubiquinone] iron-sulfur			
	protein 4, mitochondrial; Complex I-18 kDa; CI-18			
	kDa; Complex I-AQDQ; CI-AQDQ;			
	NADH-ubiquin	one oxidore	ductase 18 kDa subunit	
Immunagan	Decembinent fu	11 Ionath nu	otein of human NDUFS4	
Immunogen:	Recombinant lu	ii iengui pro	Stelli of human NDOF 54	
Form: Liquid		Concentr	ation: lmg/mL	
Size: 100 ul/50 ul		Host: Rabbit		
Reactivity: Human, Mouse, Rat		Application: WB, IF/IC		
Clonality: Polyclo			WB (1/500 - 1/2000),	




IF/IC (1/10 - 1/100)

Entrez Gene:	4724/17993/499529
SwissProt:	O43181/Q9CXZ1/Q5XIF3
Purification:	The antibody was purified by immunogen affinity chromatography.
Buffer:	Liquid in 0.42% Potassium phosphate, 0.87% Sodium chloride, pH 7.3, 30% glycerol, and 0.01% sodium azide.

WB description:

Western blot analysis of NDUFS4 expression in SW620 (A),


MCF7 (B), HepG2 (C), mouse brain (D), rat liver (E) whole cell lysates.



IF/ICC description:



Immunofluorescent analysis of NDUFS4 staining in HepG2 cells. Formalin-fixed cells were permeabilized with 0.1% Triton X-100 in TBS for 5-10 minutes and blocked with 3% BSA-PBS for 30 minutes at room temperature. Cells were probed with the primary antibod



Storage:

Store at -20°C. Avoid repeated freeze / thaw cycles.

