

| Manufacturer/Su | nnlier | | | |
|---|---|---|----------------------|--|
| WUXI DONGLIN SCI&TECH DEVELOPMENT CO., LTD | | | | |
| Address: A1-203 Mingpin CityII,No.8 Xihudong Road.Liangxi District | | | | |
| Wuxi Jiangsu Province, China | | | | |
| TELE: 86-510-82732223 | | FAX: | 86-510-82720101-8014 | |
| Web : <u>www.dldevelop.com</u> | | | | |
| Email: <u>info@dldevelop.com.cn</u> <u>service@dldevelop.com.cn</u> | | | | |
| Product name: Catalog: | Rabbit anti-PCM DL97000A | 11 Antibod | y | |
| Synonyms: | Pericentriolar m | aterial 1 pro | otein; PCM-1; hPCM-1 | |
| Immunogen: | Recombinant full length protein of human PCM1 | | | |
| Form: Liquid | | Concentra | ation: 1mg/mL | |
| Size: 100 ul/50 ul | | Host: Rabbit | | |
| Reactivity: Human, Mouse, Rat | | Application: WB, IHC | | |
| Clonality: Polyclonal | | Dilution: WB (1/500 - 1/2000), IHC (1/50 - 1/200) | | |
| Entrez Gene: | 5108/18536 | | | |


www.dldevelop.com



| SwissProt: | Q15154/Q9R0L6 |
|---------------|---|
| Purification: | The antibody was purified by immunogen affinity chromatography. |
| Buffer: | Liquid in 0.42% Potassium phosphate, 0.87% Sodium chloride, pH 7.3, 30% glycerol, and 0.01% sodium azide. |

WB description:

Western blot analysis of PCM1 expression in SW480 (A), 293T (B) whole cell lysates.



IHC description:

Immunohistochemical analysis of PCM1 staining in mouse liver formalin fixed paraffin embedded tissue section. The section was



pre-treated using heat mediated antigen retrieval with sodium citrate buffer (pH 6.0). The section was then incubated with the antibody at room temperature and detected using an HRP conjugated compact polymer system. DAB was used as the chromogen. The section was then counterstained with haematoxylin and mounted with DPX.

Storage:

Store at -20°C. Avoid repeated freeze / thaw cycles.

