

## Manufacturer/Supplier:

WUXI DONGLIN SCI&TECH DEVELOPMENT CO., LTD

Address: A1-203 Mingpin CityII, No.8 Xihudong Road. Liangxi District

Wuxi Jiangsu Province, China

TELE: 86-510-82732223 FAX: 86-510-82720101-8014

**Web**: www.dldevelop.com

Email: info@dldevelop.com.cn service@dldevelop.com.cn

Product name: Rabbit anti-TARDBP Antibody

**Catalog:** DL97130A

**Synonyms:** TDP43; TAR DNA-binding protein 43; TDP-43

Immunogen: Recombinant full length protein of human TARDBP

Form: Liquid Concentration: 1mg/mL

Host: Rabbit **Size:** 100 ul/50 ul

Reactivity: Human, Mouse, Rat Application: WB, IHC, IF/IC, IP

**Clonality:** Polyclonal **Dilution:** WB (1/500 - 1/2000),

IHC (1/50 - 1/200), IF/IC (1/20 -

1/100), IP (1/20 - 1/50)



Entrez Gene: 23435/230908

O13148/O921F2 SwissProt:

**Purification:** The antibody was purified by immunogen affinity

chromatography.


**Buffer:** Liquid in 0.42% Potassium phosphate, 0.87%

Sodium chloride, pH 7.3, 30% glycerol, and 0.01%

sodium azide.

#### **WB** description:

Western blot analysis of TARDBP expression in Jurkat (A), SW620 (B), A673 (C) whole cell lysates.



# **IHC description:**


Immunohistochemical analysis of TARDBP staining in human stomach formalin fixed paraffin embedded tissue section. The section



was pre-treated using heat mediated antigen retrieval with sodium citrate buffer (pH 6.0). The section was then incubated with the antibody at room temperature and detected using an HRP conjugated compact polymer system. DAB was used as the chromogen. The section was then counterstained with haematoxylin and mounted with DPX.

## IF/ICC description:

Immunofluorescent analysis of TARDBP staining in SW620 cells. Formalin-fixed cells were permeabilized with 0.1% Triton X-100 in TBS for 5-10 minutes and blocked with 3% BSA-PBS for 30 minutes at room temperature. Cells were probed with the primary antibod



**IP** description:

# **Storage:**

Store at -20°C. Avoid repeated freeze / thaw cycles.