

Manufacturer/Supplier:				
WUXI DONGLIN SCI&TECH DEVELOPMENT CO., LTD				
Address: A1-203 Mingpin CityII,No.8 Xihudong Road.Liangxi District				
Wuxi Jiangsu Province, China				
TELE: 86-510-82732223		FAX:	86-510-82720101-8014	
Web : <u>www.dldevelop.com</u>				
Email: <u>info@dldevelop.com.cn</u> <u>service@dldevelop.com.cn</u>				
Product name:	Rabbit anti-Thymidine Phosphorylase Antibody			
Catalog:	DL97159A			
Synonyms:	ECGF1; Thymidine phosphorylase; TP; Gliostatin;			
	Platelet-derived endothelial cell growth factor;			
	PD-ECGF; TdRPase			
	12 2001,14			
Immunogen:	Recombinant fu	ll length pro	otein of human	
ininunogen.	Recombinant full length protein of human Thymidine Phosphorylase			
	Thymame Phos	spilorylase		
Former Linuid		Concentr	ations 1ma/mI	
Form: Liquid		Concentr	ation: 1mg/mL	
Size: 100 ul/50 ul		Host: Rab	hit	
Size: 100 ul/30 ul		HUSt. Kat	on	
Reactivity: Human, Mouse, Rat		Application: WB, IF/IC		
i i i i i i i i i i i i i i i i		- ppirouti		
Clonality: Polyclonal		Dilution:	WB (1/500 - 1/2000),	
e 5				




IF/IC (1/50 - 1/200)

Entrez Gene:	1890/72962/315219
SwissProt:	P19971/Q99N42/Q5FVR2
Purification:	The antibody was purified by immunogen affinity chromatography.
Buffer:	Liquid in 0.42% Potassium phosphate, 0.87% Sodium chloride, pH 7.3, 30% glycerol, and 0.01% sodium azide.

WB description:


Western blot analysis of Thymidine Phosphorylase expression in A549 (A), THP1 (B) whole cell lysates.



IF/ICC description:



Immunofluorescent analysis of Thymidine Phosphorylase staining in A549 cells. Formalin-fixed cells were permeabilized with 0.1% Triton X-100 in TBS for 5-10 minutes and blocked with 3% BSA-PBS for 30 minutes at room temperature. Cells were probed with the



Storage:

Store at -20°C. Avoid repeated freeze / thaw cycles.

