

Manufacturer/Supplier:

WUXI DONGLIN SCI&TECH DEVELOPMENT CO., LTD

Address: A1-203 Mingpin CityII, No.8 Xihudong Road. Liangxi District

Wuxi Jiangsu Province, China

TELE: 86-510-82732223 FAX: 86-510-82720101-8014

Web: www.dldevelop.com

Email: info@dldevelop.com.cn service@dldevelop.com.cn

Product name: Rabbit anti-NEDD4 Antibody

Catalog: DL97207A

KIAA0093; NEDD4-1; E3 ubiquitin-protein ligase **Synonyms:**

> NEDD4; Cell proliferation-inducing gene 53 protein; Neural precursor cell expressed developmentally down-regulated protein 4;

NEDD-4

Immunogen: Recombinant full length protein of human NEDD4

Form: Liquid Concentration: 1mg/mL

Size: 100 u1/50 u1 Host: Rabbit

Reactivity: Human **Application:** WB, IHC



Clonality: Polyclonal **Dilution:** WB (1/500 - 1/2000),

IHC (1/50 - 1/200)

Entrez Gene: 4734

P46934 SwissProt:

Purification: The antibody was purified by immunogen affinity

chromatography.

Buffer: Liquid in 0.42% Potassium phosphate, 0.87%


Sodium chloride, pH 7.3, 30% glycerol, and 0.01%

sodium azide.

WB description:

Western blot analysis of NEDD4 expression in Hela (A), Jurkat (B), A549 (C), SKOV3 (D) whole cell lysates.





IHC description:

Immunohistochemical analysis of NEDD4 staining in human prostate cancer formalin fixed paraffin embedded tissue section. The section was pre-treated using heat mediated antigen retrieval with sodium citrate buffer (pH 6.0). The section was then incubated with the antibody at room temperature and detected using an HRP conjugated compact polymer system. DAB was used as the chromogen. The section was then counterstained with haematoxylin and mounted with DPX.

Storage:

Store at -20°C. Avoid repeated freeze / thaw cycles.