

Manufacturer/Supplier:				
WUXI DONGLIN SCI&TECH DEVELOPMENT CO., LTD				
Address: A1-203 Mingpin CityII,No.8 Xihudong Road.Liangxi District				
Wuxi Jiangsu Province, China				
TELE: 86-510-82732223 FAX: 86-510-82720101-801				
Web : <u>www.dldevelop.com</u>				
Email: info@dldevelop.com.cn service@dldevelop.com.cn				
Product name:	Rabbit anti-Histone H3 (MonoMethyl K4) Antibody			
Catalog:	DL97682A			
0				
Synonyms:	H3FT; Histone H3.1t; H3/t; H3t; H3/g			
	1101 1, 11100010		1150, 115, 5	
Immunogen:	A synthetic met	avlated non	tida corresponding to	
minunogen.	A synthetic methylated peptide corresponding to			
	residues surrounding K4 of human Histone H3			
	(MonoMethyl K	.4)		
Form: Liquid		Concentr	ation: 1mg/mL	
Size: 100 ul/50 ul		Host: Rab	bit	
Reactivity: Human, Mouse, Rat		Applicati	on: WB, IHC, IF/IC, IP,	
	, ,	CHIP))))	
Clonality: Polyclonal		01111	WB (1/500 - 1/2000),	
cronanty. i orycioliai		IHC (1/50 - 1/200), IF/IC (1/50 -		
		IHC (1/50	- 1/200), IF/IC (1/30 -	




1/200), IP (1/50 - 1/100), ChIP (1/100 - 1/500)

Entrez Gene:	8290
SwissProt:	Q16695
Purification:	The antibody was purified by immunogen affinity chromatography.
Buffer:	Liquid in 0.42% Potassium phosphate, 0.87% Sodium chloride, pH 7.3, 30% glycerol, and 0.01% sodium azide.

WB description:

Western blot analysis of Histone H3 (MonoMethyl K4) expression in Hela (A) whole cell lysates.

DEVELOP




IHC description:

Immunohistochemical analysis of Histone H3 (MonoMethyl K4) staining in human brain formalin fixed paraffin embedded tissue section. The section was pre-treated using heat mediated antigen retrieval with sodium citrate buffer (pH 6.0). The section was then incubated with the antibody at room temperature and detected using an HRP conjugated compact polymer system. DAB was used as the chromogen. The section was then counterstained with haematoxylin and mounted with DPX.

IF/ICC description:

Immunofluorescent analysis of Histone H3 (MonoMethyl K4) staining in HEK293T cells. Formalin-fixed cells were permeabilized with 0.1% Triton X-100 in TBS for 5-10 minutes and blocked with 3% BSA-PBS for 30 minutes at room temperature. Cells were probed wi

DEVELOP.



IP description:

ChIP description:

Storage:

Store at -20°C. Avoid repeated freeze / thaw cycles.

