

Manufacturer/Supplier:

WUXI DONGLIN SCI&TECH DEVELOPMENT CO., LTD

Address: A1-203 Mingpin CityII, No.8 Xihudong Road. Liangxi District

Wuxi Jiangsu Province, China

TELE: 86-510-82732223 FAX: 86-510-82720101-8014

Web: www.dldevelop.com

Email: info@dldevelop.com.cn service@dldevelop.com.cn

Product name: Rabbit anti-Histone H3 (TriMethyl K27) Antibody

Catalog: DL97688A

H3FT; Histone H3.1t; H3/t; H3t; H3/g **Synonyms:**

Immunogen: A synthetic methylated peptide corresponding to

residues surrounding K27 of human Histone H3

(TriMethyl K27)

Form: Liquid Concentration: 1mg/mL

Size: 100 ul/50 ul Host: Rabbit

Reactivity: Human, Mouse, Rat Application: WB, IHC, IF/IC, IP,

CHIP

Dilution: WB (1/500 - 1/2000), **Clonality:** Polyclonal

IHC (1/50 - 1/200), IF/IC (1/50 -



1/200), IP (1/50 - 1/100), ChIP (1/100 - 1/500)

Entrez Gene: 8290

Q16695 SwissProt:

Purification: The antibody was purified by immunogen affinity

chromatography.

Buffer: Liquid in 0.42% Potassium phosphate, 0.87%


Sodium chloride, pH 7.3, 30% glycerol, and 0.01%

sodium azide.

WB description:

Western blot analysis of Histone H3 (TriMethyl K27) expression in Hela (A) whole cell lysates.






IHC description:

Immunohistochemical analysis of Histone H3 (TriMethyl K27) staining in human brain formalin fixed paraffin embedded tissue section. The section was pre-treated using heat mediated antigen retrieval with sodium citrate buffer (pH 6.0). The section was then incubated with the antibody at room temperature and detected using an HRP conjugated compact polymer system. DAB was used as the chromogen. The section was then counterstained with haematoxylin and mounted with DPX.

IF/ICC description:

Immunofluorescent analysis of Histone H3 (TriMethyl K27) staining in HEK293T cells. Formalin-fixed cells were permeabilized with 0.1% Triton X-100 in TBS for 5-10 minutes and blocked with 3% BSA-PBS for 30 minutes at room temperature. Cells were probed wi





IP description:

ChIP description:

Storage:

Store at -20°C. Avoid repeated freeze / thaw cycles.