

Manufacturer/Supplier:				
WUXI DONGLIN SCI&TECH DEVELOPMENT CO., LTD				
Address: A1-203 Mingpin CityII,No.8 Xihudong Road.Liangxi District				
Wuxi Jiangsu Province, China				
TELE: 86-510-82732223		FAX:	86-510-82720101-8014	
Web : <u>www.dldevelop.com</u>				
Email: <u>info@dldevelop.com.cn</u> <u>service@dldevelop.com.cn</u>				
Product name:	Rabbit anti-Histone H3 (DiMethyl K79) Antibody			
Catalog:	DL97694A			
Synonyms:	H3FT; Histone H3.1t; H3/t; H3t; H3/g			
Immunogen:	KLH-conjugated synthetic peptide of human			
	Histone H3 (Dil	Methyl K79	)	
		~		
Form: Liquid		Concentra	ation: 1mg/mL	
S: 100 1/20 1		<b>H</b> ( <b>D</b> 1	1 *.	
Size: 100 ul/50 ul		Host: Rab	01	
<b>Reactivity:</b> Human, Mouse, Rat <b>Application:</b> WB, IHC, IF/IC, IP,				
Reactivity: Human, Mouse, Rat		CHIP	<b>JII.</b> WD, IIIC, II/IC, II,	
<b>Clonality:</b> Polyclonal			WB (1/500 - 1/2000),	
			- 1/200), IF/IC (1/50 -	
			(1/50 - 1/100), ChIP	
		, II	(	




## (1/100 - 1/500)

Entrez Gene:	8290
SwissProt:	Q16695
Purification:	The antibody was purified by immunogen affinity chromatography.
Buffer:	Liquid in 0.42% Potassium phosphate, 0.87% Sodium chloride, pH 7.3, 30% glycerol, and 0.01% sodium azide.

#### **WB description:**

Western blot analysis of Histone H3 (DiMethyl K79) expression in Hela (A) whole cell lysates.




### **IHC description:**



Immunohistochemical analysis of Histone H3 (DiMethyl K79) staining in human kidney formalin fixed paraffin embedded tissue section. The section was pre-treated using heat mediated antigen retrieval with sodium citrate buffer (pH 6.0). The section was then incubated with the antibody at room temperature and detected using an HRP conjugated compact polymer system. DAB was used as the chromogen. The section was then counterstained with haematoxylin and mounted with DPX.

#### **IF/ICC description:**

Immunofluorescent analysis of Histone H3 (DiMethyl K79) staining in HEK293T cells. Formalin-fixed cells were permeabilized with 0.1% Triton X-100 in TBS for 5-10 minutes and blocked with 3% BSA-PBS for 30 minutes at room temperature. Cells were probed wit



**IP description:** 

# **DEVELOP**.

## **ChIP description:**

#### **Storage:**

Store at -20°C. Avoid repeated freeze / thaw cycles.

