

## Manufacturer/Supplier:

WUXI DONGLIN SCI&TECH DEVELOPMENT CO., LTD

Address: A1-203 Mingpin CityII, No.8 Xihudong Road. Liangxi District

Wuxi Jiangsu Province, China

TELE: 86-510-82732223 FAX: 86-510-82720101-8014

Web: www.dldevelop.com

Email: <u>info@dldevelop.com.cn</u> <u>service@dldevelop.com.cn</u>

**Product name:** Rabbit anti-DNase I Antibody

Catalog: DL97706A

**Synonyms:** DNL1; DRNI; Deoxyribonuclease-1;

Deoxyribonuclease I; DNase I; Dornase alfa

Immunogen: Recombinant full length protein of human DNase I

Form: Liquid Concentration: 1mg/mL

Size: 100 ul/50 ul Host: Rabbit

Reactivity: Human, Mouse, Rat Application: WB, IHC, IF/IC

Clonality: Polyclonal Dilution: WB (1/500 - 1/2000),

IHC (1/50 - 1/200), IF/IC (1/50 -

1/200)



Entrez Gene: 1773/13419/25633

P24855/P49183/P21704 SwissProt:

Purification: The antibody was purified by immunogen affinity

chromatography.


**Buffer:** Liquid in 0.42% Potassium phosphate, 0.87%

Sodium chloride, pH 7.3, 30% glycerol, and 0.01%

sodium azide.

### **WB** description:

Western blot analysis of DNase I expression in HT29 (A), mouse testis (B) whole cell lysates.



### **IHC description:**


Immunohistochemical analysis of DNase I staining in human brain



formalin fixed paraffin embedded tissue section. The section was pre-treated using heat mediated antigen retrieval with sodium citrate buffer (pH 6.0). The section was then incubated with the antibody at room temperature and detected using an HRP conjugated compact polymer system. DAB was used as the chromogen. The section was then counterstained with haematoxylin and mounted with DPX.

# IF/ICC description:

Immunofluorescent analysis of DNase I staining in HeLa cells. Formalin-fixed cells were permeabilized with 0.1% Triton X-100 in TBS for 5-10 minutes and blocked with 3% BSA-PBS for 30 minutes at room temperature. Cells were probed with the primary antibod



### **Storage:**

Store at -20°C. Avoid repeated freeze / thaw cycles.