

Manufacturer/Supplier:

WUXI DONGLIN SCI&TECH DEVELOPMENT CO., LTD

Address: A1-203 Mingpin CityII, No.8 Xihudong Road. Liangxi District
Wuxi Jiangsu Province, China

TELE: 86-510-82732223

FAX: 86-510-82720101-8014

Web : www.dldevelop.com

Email: info@dldevelop.com.cn service@dldevelop.com.cn

Product name: Rabbit anti-ERCC2 Antibody

Catalog: DL97958A

Synonyms: XPD; XPDC; TFIIH basal transcription factor complex helicase XPD subunit; Basic transcription factor 2 80 kDa subunit; BTF2 p80; CXPB; DNA excision repair protein ERCC-2; DNA repair protein complementing XP-D cells; TFIIH basal transcription factor complex

Immunogen: Recombinant full length protein of human ERCC2

Form: Liquid

Concentration: 1mg/mL

Size: 100 ul/50 ul

Host: Rabbit

Reactivity: Human, Mouse, Rat

Application: WB, IF/IC

Clonality: Polyclonal

Dilution: WB (1/500 - 1/2000),
IF/IC (1/50 - 1/200)

Entrez Gene: 2068/13871


SwissProt: P18074/O08811

Purification: The antibody was purified by immunogen affinity chromatography.

Buffer: Liquid in 0.42% Potassium phosphate, 0.87% Sodium chloride, pH 7.3, 30% glycerol, and 0.01% sodium azide.


WB description:

Western blot analysis of ERCC2 expression in THP1 (A), HepG2 (B) whole cell lysates.



IF/ICC description:

Immunofluorescent analysis of ERCC2 staining in HeLa cells. Formalin-fixed cells were permeabilized with 0.1% Triton X-100 in TBS for 5-10 minutes and blocked with 3% BSA-PBS for 30 minutes at room temperature. Cells were probed with the primary antibody



Storage:



Store at -20°C. Avoid repeated freeze / thaw cycles.