

Manufacturer/Supplier:

WUXI DONGLIN SCI&TECH DEVELOPMENT CO., LTD

Address: A1-203 Mingpin CityII, No.8 Xihudong Road. Liangxi District

Wuxi Jiangsu Province, China

TELE: 86-510-82732223 FAX: 86-510-82720101-8014

Web: www.dldevelop.com

Email: info@dldevelop.com.cn service@dldevelop.com.cn

Product name: Rabbit anti-TRIM69 Antibody

Catalog: DL97990A

Synonyms: RNF36; E3 ubiquitin-protein ligase TRIM69;

RFP-like domain-containing protein trimless; RING finger protein 36; Tripartite motif-containing protein

69

Immunogen: Recombinant full length protein of human TRIM69

Form: Liquid Concentration: 1mg/mL

Size: 100 ul/50 ul Host: Rabbit

Reactivity: Human, Mouse, Rat **Application:** WB, IHC, IF/IC

Clonality: Polyclonal Dilution: WB (1/500 - 1/2000),



IHC (1/50 - 1/200), IF/IC (1/50 -1/200)

Entrez Gene: 140691/70928/311373

Q86WT6/Q80X56/Q5BK82 SwissProt:

Purification: The antibody was purified by immunogen affinity

chromatography.

Buffer: Liquid in 0.42% Potassium phosphate, 0.87%


Sodium chloride, pH 7.3, 30% glycerol, and 0.01%

sodium azide.

WB description:

Western blot analysis of TRIM69 expression in mouse liver (A) whole cell lysates.






IHC description:

Immunohistochemical analysis of TRIM69 staining in human testis formalin fixed paraffin embedded tissue section. The section was pre-treated using heat mediated antigen retrieval with sodium citrate buffer (pH 6.0). The section was then incubated with the antibody at room temperature and detected using an HRP conjugated compact polymer system. DAB was used as the chromogen. The section was then counterstained with haematoxylin and mounted with DPX.

IF/ICC description:

Immunofluorescent analysis of TRIM69 staining in A549 cells. Formalin-fixed cells were permeabilized with 0.1% Triton X-100 in TBS for 5-10 minutes and blocked with 3% BSA-PBS for 30 minutes at room temperature. Cells were probed with the primary antibody





Storage: Store at -20°C. Avoid repeated freeze / thaw cycles.