

Manufacturer/Supplier:

WUXI DONGLIN SCI&TECH DEVELOPMENT CO., LTD

Address: A1-203 Mingpin CityII, No.8 Xihudong Road, Liangxi District
Wuxi Jiangsu Province, China

TELE: 86-510-82732223

FAX: 86-510-82720101-8014

Web : www.dldevelop.com

Email: info@dldevelop.com.cn service@dldevelop.com.cn

Product name: Rabbit anti-eIF2C2 Antibody

Catalog: DL98081A

Synonyms: EIF2C2; Protein argonaute-2; Argonaute2; hAgo2;
Argonaute RISC catalytic component 2; Eukaryotic
translation initiation factor 2C 2; eIF-2C 2; eIF2C 2;
PAZ Piwi domain protein; PPD; Protein slicer

Immunogen: Recombinant full length protein of human eIF2C2

Form: Liquid

Concentration: 1mg/mL

Size: 100 ul/50 ul

Host: Rabbit

Reactivity: Human, Mouse, Rat

Application: WB, IHC, IP

Clonality: Polyclonal

Dilution: WB (1/500 - 1/2000),

IHC (1/50 - 1/200), IP (1/50 -
1/100)

Entrez Gene: 27161/239528/59117


SwissProt: Q9UKV8/Q8CJG0/Q9QZ81

Purification: The antibody was purified by immunogen affinity chromatography.

Buffer: Liquid in 0.42% Potassium phosphate, 0.87% Sodium chloride, pH 7.3, 30% glycerol, and 0.01% sodium azide.

WB description:

Western blot analysis of eIF2C2 expression in Jurkat (A) whole cell lysates.



IHC description:

Immunohistochemical analysis of eIF2C2 staining in human brain formalin fixed paraffin embedded tissue section. The section was pre-treated using heat mediated antigen retrieval with sodium citrate buffer (pH 6.0). The section was then incubated with the antibody at room temperature and detected using an HRP conjugated compact polymer system. DAB was used as the chromogen. The section was then counterstained with haematoxylin and mounted with DPX.

IP description:

Storage:

Store at -20°C. Avoid repeated freeze / thaw cycles.