

Manufacturer/Supplier:

WUXI DONGLIN SCI&TECH DEVELOPMENT CO., LTD

Address: A1-203 Mingpin CityII, No.8 Xihudong Road. Liangxi District

Wuxi Jiangsu Province, China

TELE: 86-510-82732223 FAX: 86-510-82720101-8014

Web: www.dldevelop.com

Email: info@dldevelop.com.cn service@dldevelop.com.cn

Product name: Rabbit anti-C8orf79 Antibody

Catalog: DL98153A

Synonyms: C8orf79; Uncharacterized protein KIAA1456

Immunogen: Recombinant full length protein of human C8orf79

Form: Liquid Concentration: 1mg/mL

Size: 100 ul/50 ul Host: Rabbit

Reactivity: Human, Mouse, Rat Application: WB, IHC, IF/IC

Clonality: Polyclonal Dilution: WB (1/500 - 1/2000),

IHC (1/50 - 1/200), IF/IC (1/50 -

1/200)



Entrez Gene: 57604

Q8N9K7 SwissProt:

Purification: The antibody was purified by immunogen affinity

chromatography.


Buffer: Liquid in 0.42% Potassium phosphate, 0.87%

Sodium chloride, pH 7.3, 30% glycerol, and 0.01%

sodium azide.

WB description:

Western blot analysis of C8orf79 expression in HepG2 (A), mouse pancreas (B) whole cell lysates.



IHC description:


Immunohistochemical analysis of C8orf79 staining in human prostate cancer formalin fixed paraffin embedded tissue section. The



section was pre-treated using heat mediated antigen retrieval with sodium citrate buffer (pH 6.0). The section was then incubated with the antibody at room temperature and detected using an HRP conjugated compact polymer system. DAB was used as the chromogen. The section was then counterstained with haematoxylin and mounted with DPX.

IF/ICC description:

Immunofluorescent analysis of C8orf79 staining in A549 cells. Formalin-fixed cells were permeabilized with 0.1% Triton X-100 in TBS for 5-10 minutes and blocked with 3% BSA-PBS for 30 minutes at room temperature. Cells were probed with the primary antibod



Storage:

Store at -20°C. Avoid repeated freeze / thaw cycles.